

Violation of Leggett-Garg inequalities in single quantum dot

Yong-Nan Sun, Yang Zou, Rong-Chun Ge, Jian-Shun Tang, Chuan-Feng Li *, and Guang-Can Guo

*Key Laboratory of Quantum Information, University of Science and Technology of China,
CAS, Hefei, 230026, People's Republic of China*

(Dated: October 27, 2011)

We investigate the violation of Leggett-Garg (LG) inequalities in quantum dots with the stationarity assumption. By comparing two types of LG inequalities, we find a better one which is easier to be tested in experiment. In addition, we show that the fine-structure splitting, background noise and temperature of quantum dots all influence the violation of LG inequalities.

PACS numbers: 03.67.Mn, 42.50.-p, 78.67.Hc

The transition from the strange quantum description to our familiar classical description is one of the fundamental questions in understanding the world. This issue was first investigated by Leggett and Garg in 1985 [1] and led to the formulation of the so called temporal Bell inequalities. Instead of the correlations between states of two spatially separated system, they cared about the correlations of the state of a single system at different time. What's more, in order to get the inequalities, Leggett and Garg made two general assumptions [2]: (1) any two-level macroscopic system will be at any time in one of the two accessible states (Macroscopic realism). (2) the actual state of the system can be determined with arbitrarily small perturbation of its subsequent dynamics (Non-invasive measurability). Leggett-Garg (LG) inequalities provide a criterion to characterize the boundary between the quantum realm and classical one and the possibility of identifying the macroscopic quantum coherence.

Due to the assumption of noninvasive measurement, which describes the ability to determine the state of the system without any influence on its subsequent dynamics, it is difficult to test the LG inequalities experimentally. There have been many proposals to test such kind of inequalities by employing superconducting quantum interference devices [1, 3, 4], but the extreme difficulty of experiments with truly macroscopic systems, as well as the request of noninvasive measurement, has made it impossible to draw any clear conclusion. The LG inequalities have been tested in optical system with CNOT gate [5]. A more recent approach takes advantage of weak measurement [6–8], in which the dynamics of the system is slightly disturbed. In this work, we follow a different approach. Testable LG inequalities is derived by replacing the noninvasive measurement assumption with the stationarity assumption [9, 10].

Considering an observable $Q(t)$ of a two level-system such as the polarization of the photon, in this case, we can define $Q(t) = 1$ when the state of the photon is $|H\rangle$, and $Q(t) = -1$ when the state of the photon is $|V\rangle$. The autocorrelation function of this observable is defined as $K(t_1, t_2) = \langle Q(t_1)Q(t_2) \rangle$. For three different times t_1 ,

t_2 and t_3 ($t_1 < t_2 < t_3$), we get the two LG inequalities under the realistic description [1]

$$K(t_1, t_2) + K(t_2, t_3) + K(t_1, t_3) \geq -1, \quad (1)$$

$$K(t_1, t_3) - K(t_1, t_2) - K(t_2, t_3) \geq -1. \quad (2)$$

Once the assumption of stationarity is introduced, in the case of $t_2 - t_1 = t_3 - t_2 = t$, the inequalities become [9]

$$K_+ = K(2t) + 2K(t) \geq -1, \quad (3)$$

$$K_- = K(2t) - 2K(t) \geq -1. \quad (4)$$

These inequalities set the boundary of the temporal correlations and are amenable to an experimental testing by a two-shot experiment.

In this paper, we compare the two types of LG inequalities and discuss the violation of LG inequalities with the influence of the fine-structure splitting, background noise and temperature in a quantum dot system.

Semiconductor quantum dots, often referred to as “artificial atoms”, have well defined discrete energy levels [11] due to their three-dimensional confinement of electrons. The atomlike properties of the single semiconductor quantum dot, together with its ease of integration into more complicated device structures, have made it an attractive and widely-studied system for applications in quantum information [12]. One of their potentially useful properties is the emission of polarization entangled photon pairs by the radiative decay of the biexciton state [13–16].

The energy level of a quantum dot is shown in Figure 1, we care about the two exciton states $|2\rangle$ and $|1\rangle$ in a quantum dot. We define $Q(t) = 1$ when the quantum dot state is $\frac{1}{\sqrt{2}}(|2\rangle + |1\rangle)$ and $Q(t) = -1$ when the quantum dot state is $\frac{1}{\sqrt{2}}(|2\rangle - |1\rangle)$. The quantum dot is initially excited to the biexciton state by a short-pulsed laser and then evolves in a thermal bath. After emitting a biexciton photon, the proposed single quantum dot system will be in an entangled photon-exciton state. For an ideal quantum dot with degenerate intermediate exciton states $|2\rangle$ and $|1\rangle$, it is a maximum entangled state $|\psi\rangle = \frac{1}{\sqrt{2}}[|H\rangle|2\rangle + |V\rangle|1\rangle]$. Then if we find the biexciton

*email: cfli@ustc.edu.cn




FIG. 1: (Color online). Energy-level schematic of the biexciton cascade process. The ground state $|0\rangle$, the two linear-polarized exciton state $|2\rangle$ and $|1\rangle$, the biexciton state $|3\rangle$, the fine-structure splitting S .

photon in state $\frac{1}{\sqrt{2}}(|H\rangle + |V\rangle)$, the state of the quantum dot system will be $\frac{1}{\sqrt{2}}(|2\rangle + |1\rangle)$. With the help of measurement, we can prepare the initial state of the quantum dot system. After a period of time of evolution, the quantum dot system will emit the second exciton photon. Then, if the state of the second photon is detected to be $\frac{1}{\sqrt{2}}(|H\rangle - |V\rangle)$, it indicates the quantum dot system ends up in $\frac{1}{\sqrt{2}}(|2\rangle - |1\rangle)$. In this case, because of the relationship between photon and the quantum dot system, we can detect the photon state to evaluate the LG inequalities. With these photon states, we can evaluate the LG inequalities.

Taking the acoustic phonon-assisted transition process into account, the density matrix of this four-level system can be described with the master equation [17]

$$\dot{\hat{\rho}} = -i[\hat{H}_0, \hat{\rho}] + L(\hat{\rho}), \quad (5)$$

where $\hat{H}_0 = \sum_{i=0}^3 \omega_i |i\rangle\langle i|$ and $L(\hat{\rho})$ is the Lindblad term denotes the dissipation term [18]. When projected onto the subspace spanned by the four basis $\{|H_1 H_2\rangle, |H_1 V_2\rangle, |V_1 H_2\rangle, |V_1 V_2\rangle\}$, we can get the corresponding two-photon polarization density matrix $\hat{\rho}_{pol}$. With the presence of the background noise and the overlap of the photon pairs frequency distributions, the total density matrix of the photon pairs is divided into three parts [18]

$$\hat{\rho}_{tot} = \frac{1}{1+g} [\eta \hat{\rho}_{pol} + (1-\eta) \hat{\rho}_{noc} + g \hat{\rho}_{noise}], \quad (6)$$

where $\hat{\rho}_{noc}$ arises from the fine-structure splitting induced distinguishability between the two decay paths, representing the nonoverlap of the photon frequency distributions. The third term $\hat{\rho}_{noise}$, which describes the background noise, is set as an identity matrix. At last, the

total density matrix of the photon pairs can be expressed as follows

$$\hat{\rho}_{tot} = \begin{pmatrix} \rho_{11} & 0 & 0 & \rho_{14} \\ 0 & \rho_{22} & 0 & 0 \\ 0 & 0 & \rho_{33} & 0 \\ \rho_{41} & 0 & 0 & \rho_{44} \end{pmatrix}$$

In our model, we can define the observable $Q(t) = 1$ when the state of the photon is $|+\rangle = \frac{1}{\sqrt{2}}(|H\rangle + |V\rangle)$, and $Q(t) = -1$ when the state of the photon is $|-\rangle = \frac{1}{\sqrt{2}}(|H\rangle - |V\rangle)$. At the time $t = 0$, we detect the first photon under the bases $|+\rangle$ and postselect the result of $|+\rangle$ to prepare the initial state of the quantum dot system. After a period of time t , we detect the second photon under the bases $|+\rangle$ and $|-\rangle$. We define the joint probability P_{++} when the state of the second photon is $|+\rangle$ and the probability P_{+-} when the state of the second photon is $|-\rangle$. We detect the photon pairs with delays τ in the range $t \leq \tau \leq (t+\omega)$, by employing a single timing gate [19], where t is the start time of the gate. At last, we can evaluate the autocorrelation by the expression

$$K(t) = P_{++}(t) - P_{+-}(t). \quad (7)$$

The autocorrelation will be measured two times by two independent experiments which begin with a primary system in an identical initial state and evolves under identical conditions. In the first experiment, the measurement of the autocorrelation is made at the time t and the second measurement is made at the time $2t$ by another experiment and the joint probability can be calculated through the total density matrix of the photon pairs, then the final autocorrelation can be expressed as follows

$$K(t) = \frac{\rho_{14}(t) + \rho_{41}(t)}{\rho_{11}(t) + \rho_{22}(t) + \rho_{33}(t) + \rho_{44}(t)}, \quad (8)$$

Therefore, we can evaluate the LG inequalities through the expression $K_+ = K(2t) + 2K(t)$, and $K_- = K(2t) - 2K(t)$.

Our result is shown in Fig. 2. We can see that the curve of the LG oscillates with time, what's more, the amplitude of the curve decays when the time increases. By comparing the two types of LG inequalities, we find that the K_- reaches the classical limit -1 first. As we know, when the time delay of the photon pairs increase, the biphoton coincidence decreases. Therefore, K_- is easier to be tested in the experiment and we discuss K_- for the following results.

Then we analysis the relationship between the LG inequality and the fine-structure splitting. The result is presented in Fig. 3. In this case, the background noise $g = 0.3$, the gate width $\omega = 50$ ps and the temperature $T = 5$ K. When the fine-structure splitting is small, the LG inequality is violated easily. However, with the increase of the fine-structure splitting, the violation of LG inequality becomes increasingly weak. It can be seen that




FIG. 2: Two types of LG inequalities. The fine-structure splitting $S = 3 \mu\text{eV}$, background noise $g=0$, the gate width $\omega = 50 \text{ ps}$ and the temperature $T = 5 \text{ K}$.




FIG. 3: The relationship between fine-structure splitting and Leggett-Garg inequality. In the conditions of the background noise $g = 0.3$, the gate width $\omega = 50 \text{ ps}$ and the temperature $T = 5 \text{ K}$. The violation of the LG inequality becomes easier when the fine-structure splitting decreases.

K_- do not violate the classical limit -1 when the fine-structure splitting becomes large enough. This implies that when the fine-structure splitting becomes large, the evolution process can be described by classical realistic theory.

Further more, the influence of background noise and




FIG. 4: The relationship between background noise and the LG inequality. The fine-structure splitting $S = 3 \mu\text{eV}$, the gate width $\omega = 50 \text{ ps}$ and the temperature $T = 5 \text{ K}$.




FIG. 5: The relationship between temperature and the LG inequality. The fine-structure splitting $S = 2.5 \mu\text{eV}$, the background noise $g = 0.2$ and the gate width $\omega = 50 \text{ ps}$.

temperature are investigated. As shown in Fig. 4 and Fig. 5, when the background noise and temperature decrease, the curve of the LG inequality goes below the classical limit -1 easily. The transition from quantum process to classical process is also been seen clearly.

At last, it should be noticed that there is an approximation in our model. We use the state of photon emitted to determine the state of quantum dot, this is true when the fine structure splitting is zero. In real situation, we need the fine structure splitting to be nonzero in order to introduce the evolution, the partial distinguishability of spectrums between $|H\rangle$ and $|V\rangle$ will introduce errors. However, it can be easily verified that the errors only make the LG inequality more difficult to be violated, i.e., the requirement is not loosen in our model.

In summary, we have investigated the violation of the LG inequalities in quantum dot system. When the fine-structure splitting of the quantum dot system, background noise and the temperature become small, we achieve the maximal violation. Better results may be obtained, when we change the method by which we get the initial state. Finally, we can see clearly from the results that the LG inequalities can be used as a criterion to identify the boundary of the classical realistic description.

This work was supported by National Fundamental Research Program, National Natural Science Foundation of China (Grant Nos. 60921091 and 10874162).

-
- [1] A. J. Leggett and A. Garg, Phys. Rev. Lett. 54, 857 (1985).
 - [2] S. F. Huelga, T. W. Marshall, and E. Santos, Europhys. Lett. 38, 249 (1997)
 - [3] C. D. Tesche, Phys. Rev. Lett. 64, 2358 (1990).
 - [4] J. P. Paz and G. Mahler, Phys. Rev. Lett. 71, 3235 (1993).
 - [5] J. S. Xu, C. F. Li, X. B. Zou, and G. C. Guo, Sci. Rep. 1, 101 (2011).
 - [6] A. Palacios-Laloy, F. Mallet, F. Nguyen, P. Bertet, D. Vion, D. Esteve, and A. N. Korotov, Nature Phys. 6, 442 (2010).
 - [7] J. Dressel, C. J. Broadbent, J. C. Howell, and A. N. Jordan, Phys. Rev. Lett. 106, 040402 (2011).
 - [8] M. E. Goggin, M. P. Almeida, M. Barbieri, B. P. Lanyon, J. L. O'Brien, A. G. White, and G. J. Pryde, Proc. Natl. Acad. Sci. U.S.A. 108, 1256 (2011).
 - [9] S. F. Huelga, T. W. Marshall, and E. Santos, Phys. Rev. A 52, R2497 (1995).
 - [10] S. F. Huelga, T. W. Marshall, and E. Santos, Phys. Rev. A 54, 1798 (1996).
 - [11] A. J. Shields, Nat. Photon. 1, 215 (2007).
 - [12] Daniel Loss and David P. Divincenzo, Phys. Rev. A 57, 120 (1998).
 - [13] O. Benson, C. Santori, M. Pelton, and Y. Yamamoto, Phys. Rev. Lett. 84, 2513 (2000).
 - [14] R. M. Stevenson, R. J. Young, P. Atkinson, K. Cooper, D. A. Ritchie, A. J. Shields, Nature (London) 439, 179 (2006).
 - [15] R. J. Young, R. M. Stevenson, P. Atkinson, K. Cooper, D. A. Ritchie, A. J. Shields, New J. Phys. 8, 29 (2006).
 - [16] R. M. Stevenson, A. J. Hudson, R. J. Young, P. Atkinson, K. Cooper, D. A. Ritchie, A. J. Shields, Opt. Express 15, 6507 (2007).
 - [17] Q. Q. Wang, A. Muller, M. T. Cheng, H. J. Zhou, P. Bianucci, and C. K. Shih, Phys. Rev. Lett. 95, 187404 (2005).
 - [18] Y. Zou, M. Gong, C. F. Li, G. Chen, J. S. Tang, and G. C. Guo, Phys. Rev. A 81, 064303 (2010).
 - [19] R. M. Stevenson, A. J. Hudson, A. J. Bennett, R. J. Young, C. A. Nicoll, D. A. Ritchie, A. J. Shields, Phys. Rev. Lett 101, 170501 (2008).